

# The Clinical Significance of Posterior Insular Volume in Adolescent Anorexia Nervosa

Nancy L. Zucker, PhD, Philip A. Kragel, PhD, Henry Ryan Wagner, PhD, Lori Keeling, MA, Emeran Mayer, MD, Joyce Wang, BS, Min Su Kang, BS, Rhonda Merwin, PhD, W. Kyle Simmons, PhD, and Kevin S. LaBar, PhD

## ABSTRACT

**Objective:** The diagnostic criterion *disturbance in the experience of the body* remains a poorly understood and persistent feature of anorexia nervosa (AN). Increased sophistication in understanding the structure of the insular cortex—a neural structure that receives and integrates visceral sensations with action and meaning—may elucidate the nature of this disturbance. We explored age, weight status, illness severity, and self-reported body dissatisfaction associations with insular cortex volume.

**Methods:** Structural magnetic resonance imaging data were collected from 21 adolescents with a history of AN and 20 age-, sex-, and body mass index–matched controls. Insular cortical volumes (bilateral anterior and posterior regions) were identified using manual tracing.

**Results:** Volumes of the right posterior insula demonstrated the following: (a) a significant age by clinical status interaction ( $\beta = -0.018$  [0.008];  $t = 2.32$ ,  $p = .02$ ) and (b) larger volumes were associated with longer duration of illness ( $r = 0.48$ ,  $p < .04$ ). In contrast, smaller volumes of the right anterior insula were associated with longer duration of illness ( $r = -0.50$ ,  $p < .03$ ). The associations of insular volume with body dissatisfaction were of moderate effect size and also of opposite direction, but a statistical trend in right posterior ( $r = 0.40$ ,  $p < .10$  in right posterior;  $r = -0.49$ ,  $p < .04$  in right anterior).

**Conclusions:** In this exploratory study, findings of atypical structure of the right posterior insular cortex point to the importance of future work investigating the role of visceral afferent signaling in understanding disturbance in body experience in AN.

**Key words:** interoception, anorexia nervosa, insula, body image disturbance, visceral hypersensitivity.

## INTRODUCTION

Part of the core phenomenology of anorexia nervosa (AN) is that of having one's attention intrusively captured by the experience of one's body, combined with the presence of dangerous weight loss behaviors that result in the suppression of or alteration in somatic experiences (e.g., bradycardia, bradygastria, and reduced hormonal surges with menstruation) (1,2). The elevated mortality associated with AN is well documented: AN remains one of the leading causes of premature mortality due to psychiatric causes with a standardized mortality ratio of 5.9 (95% confidence interval = 4.2–8.3) and a standardized mortality ratio attributable to suicide of 31 (95% confidence interval = 21–44) relative to other forms of mental illness or population controls (3). Although there have been impressive advances in the treatment of AN, particularly for adolescents (4–6), treatments for both adults and adolescents have had limited effectiveness in improving disturbance in the experience of the body (often referred to as body image disturbance) (1,7,8). The degree of body image disturbance predicts poor treatment response (9) and has been reported to motivate hazardous weight loss behaviors (1,2). Elucidation of the biological

substrate of body image disturbance is thus considered essential to understanding the pathophysiology of AN.

Body image disturbance is a complex construct comprising cognitive (e.g., preoccupation with appearance), attitudinal (e.g., evaluative judgments about weight), perceptual (e.g., ability to perceive body size), and experiential components (e.g., irregularities in interoceptive sensitivity). Although the cognitive, evaluative, and, increasingly, perceptual aspects of body image disturbance are well characterized (e.g., Refs. (10,11), there has been far less study of the experiential components of body image disturbance (12). Accumulating evidence suggests that aberrations in body experience in AN extend well beyond the restricted domain of weight and shape (as specified in the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition* (13)) and may include aberrations in basic somatosensory and interoceptive experiences (e.g., of heartbeat detection and tactile discrimination) as indexed by objective laboratory tasks (see review by Gaudio et al.) (12,14–16). Individuals with AN also report more intrusive sensory and interoceptive symptoms

AN = anorexia nervosa, BMI = body mass index, HC = healthy control, MRI = magnetic resonance imaging

## SDC Supplemental Content

From the Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine (Zucker, Wagner, Merwin, LaBar), Durham, North Carolina; Department of Psychology and Neuroscience, Duke University (Zucker, Kragel, Keeling, Wang, Kang, LaBar), Durham, North Carolina; Departments of Medicine, Physiology, Psychiatry & Biobehavioral Sciences, David Geffen School of Medicine, UCLA (Mayer), Los Angeles, California; and Laureate Institute, University of Tulsa (Simmons), Tulsa, Oklahoma.

Address correspondence and reprint requests to Nancy L. Zucker, PhD, Department of Psychiatry and Behavioral Science, Duke University School of Medicine, PO Box 3454, Durham, NC 27710. E-mail: zucke001@mc.duke.edu

Received for publication April 3, 2016; revision received April 2, 2017.

DOI: 10.1097/PSY.0000000000000510

Copyright © 2017 by the American Psychosomatic Society

whether currently ill or weight restored relative to healthy controls, intimating that they experience such signals as aberrant (17). This somatic sensitivity was positively associated with an evaluative measure of body dissatisfaction, suggesting that interoceptive signals and more top-down evaluative judgments of the body might be interrelated and that dietary restriction was, in part, motivated by a desire to suppress sensory sensitivity (17,18).

Why interoceptive signals and negative attitudes about the body may be associated is unclear, but some common features of AN, such as “feeling fat,” offer a hypothetical bridge as to how interoceptive experience, interpretation of that experience, and actions motivated by that experience may link these concepts. To clarify, there is evidence that those who later develop AN have increased experiences of early life events that may increase sensitivity to and preoccupation with somatic experience. For example, there is evidence of an excess of early gastrointestinal events, eating difficulties, and somatic symptoms before illness onset in AN, with sufficient evidence supporting digestive problems as a risk factor for AN (19–22). Although these early events may increase sensitization of afferent visceral pathways independently (23), such sensitivity may be combined with a learning history in which those with AN value the suppression of bodily signals or disregard the informative value of somatic signals rather than use these somatic markers to guide adaptive decision-making (24) (for review, see Ref. (25)). In the aforementioned instance, extreme sensitivity to afferent input of the gut may differentially direct attention to the body contributing to preoccupation (17), whereas learning and other environmentally influenced processes shape interpretation and meaning—including a generalized distrust or shame of bodily sensations that intrude upon awareness. In fact, Herbert et al. (26) demonstrated this dissociation in a study of intuitive eating, finding that interoceptive sensitivity and the appraisal of interoceptive signals were independently associated with the ability to eat in response to interoceptive signaling and body mass (26). Thus, one hypothesis about the persistence of body image disturbance in AN may be that the components of body image that are influenced by learning and environmental factors (e.g., interpretations and attitudes) can be addressed via treatment, whereas basic aberrations in afferent signaling from the periphery are trait features that individuals must learn to manage. However, research to date cannot distinguish whether it is something about the nature of the afferent input that is aberrant (i.e., a “bottom-up” process), a “top-down” error related to the interpretation or contextualization of somatic signals, or both.

### Insular Cortex Organization

The potential involvement of visceral sensory aberrations in AN suggests that the insular cortex is a candidate for understanding the biological basis of AN (27). The insular cortex has been described as a “center of awareness” (28) and is situated as a relay station between afferent inputs from the viscera to neural structures that help code the salience and meaning of visceral inputs (28). There have been several proposed functional mappings of the insula cortex on the basis of differences in structural architecture: a granular insula located in the posterior region, an agranular insula in the anterior region, and a dysgranular band in the middle insula (e.g., Ref. (28)); or meta-analysis (e.g., Ref. (29)); or the synthesis of resting-state connectivity with meta-analytic findings

from human neuroimaging studies (30). Consistent across these models is the delineation of a posterior insular cortex structure that is functionally connected to the somatosensory and supplementary motor areas and is associated with activities in somatosensory, sensorimotor, and pain domains (30). These inputs are then relayed forward to the anterior regions of the insular cortex in which a ventral anterior region, primarily functionally connected with limbic areas (e.g., amygdala, posterolateral orbitofrontal cortex, and ventral tegmental area), is broadly implicated in the experience and contextualization of emotional, autonomic, and chemosensory experiences (23,31). The dorsal anterior region is functionally connected to the anterior cingulate cortex and dorsolateral prefrontal cortex and is implicated in executive control and higher cognition (30). Notably, the insular cortex is one of the most widely activated neural regions across studies of awareness, emotional experience, craving/urges, and more recently, decision making (for an elegant review, see Ref. (32)). Despite this diversity in function, Craig (28) suggests that this is due to the influence of the core function of the insular cortex as an integrative site for autonomic, cognitive, and affective processing.

From a developmental standpoint, interoceptive signals initially reaching the posterior insular cortex become more elaborately deciphered and contextualized throughout maturity to increasingly guide complex decisions in both adaptive and maladaptive contexts. According to this functional neuroanatomical conceptualization, aberrant somatic experience in AN at the level of afferent input should be reflected in atypical morphology in posterior regions of the insular cortex, whereas aberrant somatic experiences at the level of interpretation and meaning would reflect atypical morphology in more anterior regions.

On the basis of the available data, it is therefore reasonable to hypothesize that individuals with AN would present with structural abnormalities in both the posterior and the anterior insula. About the directionality of hypothesized structural abnormalities, several processes may contribute to structural differences. First, early life events and premorbid activity patterns putatively support the hypothesis of increased posterior volume in AN. Early gastrointestinal pain experiences and the performance of excessive exercise or activities more generally have been documented premorbidly in AN, two classes of experience that have been associated with modulation of brain-derived neurotrophic factor, a growth factor implicated in neuronal growth, development, and survival (33,34). Speculatively, these experiences may contribute to increased neuronal growth in regions that receive somatic motor information such as the posterior insular cortex. Second, AN emerges during a peak period of brain development in which synaptic pruning contributes to the efficiency of neural networks (35). The onset of starvation during this vulnerable period may modulate normative process, with increased volume being a consequence of reduced pruning (36). Notably, a study of age-dependent changes in subcortical structure across adolescence revealed that structures such as the thalamus, implicated in somatosensory experience, exhibited an elongated developmental course relative to the cortical plate (37). Ancient cortical structures such as the insula may likewise be vulnerable to starvation if the development of the structure is elongated, particularly as the posterior portion receives somatic inputs from the thalamus and is implicated in somatosensory function (38). In a related vein, Dennis et al. (39) reported that functional connectivity of the posterior insula increased across

adolescence with temporal regions but decreased in anterior regions. Then, arguably, increased volumes in posterior insula might reflect less pruning in “early” sensory pathways, delays in neurodevelopment, or some other developmental disruption.

Given the covariation of structure and function, atypical structure at the level of the posterior insula could impact anterior insula function in that the “feed forward” nature of visceral information conveyed from the posterior to the anterior could support that misinterpretation of somatic signals in AN is based in part, on the use of aberrant sensory data in guiding behavior (40). This finding would be an important step toward advancing our understanding of the nature of body experience in AN: rather than a reflection of cognitive biases or extreme negative evaluations, body image disturbance may be a complex interplay of aberrant sensory information that influences cognitive/attitudinal factors.

### Insular Structure in AN

To date, study of insular structure in adolescent AN has been limited. In a whole-brain analysis comparing 19 adolescents with a diagnosis of AN (mean age of 15.4 years) with healthy controls, Frank et al. (41) reported increased volume in the right insular region of cases relative to controls, although insular subregions were not specified. Several recent studies that investigated structural abnormalities not restricted to the insula have been inconclusive; although focal reductions in gray matter volume (42–46) and more recently global cortical thinning (47) have been reported, other data suggest that volume reductions are secondary to starvation and remit with weight restoration (45,47). The latter findings would seem to emphasize the importance of the findings by Frank et al. (41), suggesting that increased insular volume in AN may be a vulnerability factor for the disorder.

To date, no neurobiological markers of illness severity in AN have been established, and although markers of body image disturbance are accumulating, these, too, remain limited (42,48). Thus, we undertook the current research as an exploratory study to further probe the clinical significance of insular structure in relation to clinical parameters of illness severity and to conduct exploratory analyses of insular structure in association with subjective measures of body image disturbance. We studied all adolescents with AN who presented to a specialized outpatient center for the treatment of eating disorders who either currently or previously met the criteria for AN. Each individual was characterized in terms of length of their illness and current severity of weight and attitudinal symptoms (all described hereinafter). With this sample, we addressed the following questions: (a) *Do age-related differences in the size of insular cortex regions differ in participants with a diagnosis of AN relative to sex and age-matched controls?* Our interest in age-related changes was based on the following. First, AN has a rather unique illness course, with incidence rare before pubertal onset and in adulthood. Thus, age-specific differences were of interest to inform illness pathophysiology. Second, given developmental alterations in brain structure (e.g., Refs. (35,37,49,50)), controlling for age is necessary to avoid study confounds. Although there is no evidence on the basis of clinical presentation, structural, or functional data to suspect that aberrations in body image experience would be unilateral, existing findings on insular function in AN have demonstrated right-sided aberrations. Consistent with this, we hypothesized that anatomical differences will be

right sided and will occur in both the anterior and posterior regions relative to healthy controls. Such findings would suggest differences at the level of encoding basic somatosensory input as putatively contributing to disturbances in body experience. (b) *Is volume in particular insular regions associated with duration of illness (independent of weight status or lowest BMI)?* We hypothesized that volume size in both the right anterior and right posterior insula correlates with illness parameters independent of body mass index (BMI). Associations between insular volume and illness parameters (controlling for the effects of body weight) would provide important support for the role of structural abnormalities in the insular cortex underlying the pathology in AN or on the maintenance of the disease, respectively. (c) *Will attitudinal measures of body dissatisfaction be associated with insular volume?* We hypothesized that the volume of the right posterior insular cortex would be associated with body dissatisfaction, a finding that would support the role of visceral afferent input as contributing to body image disturbance in AN.

## METHODS

### Overview

These data are part of a larger study of interoception across typical adolescence and interoception across adolescence in AN. Adolescents between the ages of 11 and 19 years were recruited for a study of “gut feelings.” Recruited individuals participated in a laboratory session and completed two magnetic resonance imaging (MRI) scans tapping decision-making and emotional regulatory capacities (reported elsewhere, see Ref. (51) and for details about these MRI tasks (52)). Here we focus on the clinical sample and report on the structural data from these scans.

### Recruitment

We attempted to recruit all individuals with AN who presented to a specialized outpatient medical clinic for the treatment of eating disorders at a Southeastern academic medical center. The control sample was recruited from this same medical clinic. Additional recruitment of both the clinical and healthy control samples was conducted throughout the university associated with this medical center.

### Healthy Control Recruitment

To maximize the generalizability of our control sample, population-based screening methods were used to screen all eligible adolescents who presented for a sick- or well-child visit to the pediatric primary care practices of a Southeastern academic medical center (see Franz et al. (53) for general screening strategy adapted for this study). The demographic composition of the primary care practice selected mimicked that of the surrounding county, thus helping to ensure the representativeness of our control group. Healthy controls were screened on random weekdays (see Fig. S1, Supplemental Digital Content 1, <http://links.lww.com/PSYMED/A410>, for study flow). Additional recruitment occurred via the student health center, counseling center, and university message boards.

### Clinical Recruitment

Clinical participants were also recruited within that medical practice. Recruiters attended every clinic session of pediatricians who were part of a specialized outpatient eating disorder program from the period of September 1, 2009, to August 31, 2011, to identify and screen all eligible AN participants, whether they currently met the criteria for AN or had a history of AN and were attending a medical follow-up appointment. Additional recruitment occurred via the student health center, counseling center, and university message boards.

### Inclusion Criteria

Clinical participants were required to have a current or previous diagnosis of AN consistent with symptoms delineated in the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition* (13). To maximize the generalizability of our sample, we were inclusive of psychiatric comorbidities. Medications were also permitted provided that the individual was on a stable dose for a period of 3 months. See Table S1, Supplemental Digital Content 1 (<http://links.lww.com/PSYMED/A410>), for medication list.

### Exclusion Criteria

Adolescents were excluded if either they or their mother did not have fluency in English, had an IQ of <70, failed to meet MRI safety requirements (see Supplemental Methods, Supplemental Digital Content 1, <http://links.lww.com/PSYMED/A410>), were suicidal, exhibited symptoms of psychosis, or actively abused substances. Healthy control participants could not have a history of an eating disorder or currently meet the criteria for a psychiatric diagnosis as determined by screening for current symptoms (see hereinafter), parent and participant report, and medical chart abstraction.

## Procedures

### Overview

Adolescents and their parents attended an initial laboratory session during which diagnostic information was obtained. The adolescent participated in a mock scanning session to familiarize herself with the scanning environment and train her in procedures that would maximize the amount of usable scanning data (e.g., teaching to minimize movement). Height and weight were obtained at the time of scanning. Individuals on medications with short half-lives (e.g., stimulant medication) were asked to refrain from taking medication on the day of scanning.

### Consent

Written informed consent was obtained from parents and participants older than 18 years; assent was obtained from participants from age 11 up to 19 years. The study was approved by the institutional review board at Duke University Medical Center (Pro00019295).

## Assessments

### Screening

Control participants were screened for the absence of mental health symptoms using questions used to predict diagnostic status from a previous population cohort study of child and adolescent psychopathology (53). Children who scored above the screen cutoff were excluded from further participation but were given a small prize.

### Determination of Diagnosis and Diagnostic History

We attempted to describe individuals with AN both categorically and dimensionally. For both, diagnosis and parameters of illness history were determined by systematically combining several sources of data: (a) maternal report of her child's illness history, (b) adolescent completion of self-report measures of current symptoms, (c) adolescent report of illness history, and (d) medical chart abstraction. This included both BMI and zBMI (i.e., age-adjusted BMI, which accounts for height, weight, and age). Categorically, groups were classified into current AN, weight-restored AN, and healthy controls (HCs). To be classified as an HC, (a) parent report indicated no history of an eating disorder, (b) adolescent self-report of Drive for Thinness was within 1 SD of normative values, and (c) the medical chart contained no reference to an eating disorder diagnosis. For an individual to be considered weight-restored AN, (a) the zBMI score had to be within 1 SD of normative values, (b) the parent report indicated that the child was without an eating disorder for 3 to 6 months, (c) the medical chart review did not contain a current diagnosis of AN, and 4) there was no evidence of a medical sign that weight

was low (e.g., bradycardia and orthostatic hypotension). For individuals to be considered AN, (a) the zBMI score was 1 SD below normative values, (b) parent records indicated the child had AN within the past 3 to 6 months, (c) medical chart had a diagnosis of AN, and (d) the adolescent had a Drive for Thinness score higher than 1 SD above normative values. To determine the length of illness, mothers were asked the age at which their child first developed an eating disorder, the type of eating disorder, and whether this diagnosis was verified by a health care professional. This information was compared and combined with the medical record and referenced against the child's weight history, current weight, and current symptom endorsement. In only one case was there a discrepancy. A similar strategy was used to determine months of weight restoration. Parents were asked the length of time the child had been at a healthy weight, and this was verified relative to the child's weight history and medical record. Again, there was one discrepancy, noted hereinafter.

### Self-Report Measures

The Eating Disorder Inventory (Third Edition) is one of the most widely used measures of eating disorder symptoms and associated features (54). This measure was used to characterize the sample relative to other studies and provide a continuous index of current symptoms. Three subscales that measure the core pathology of eating disorders were administered in the current sample: Drive for Thinness, Bulimia, and Body Dissatisfaction. All scales have extensive validity and reliability information as well as normative data from clinical and nonclinical samples. The Drive for Thinness Scale is a seven-item scale that assesses "an extreme desire to be thinner, preoccupation with weight, and an intense fear of weight gain" (pp. 14, 55). Extensive reliability, construct, and predictive validity have been established (55,56). The internal consistency of this scale in our sample was 0.95. The Bulimia subscale is an eight-item scale used to index the tendency to think about or engage in uncontrollable overeating or eating in response to emotions. Along with demonstrations of discriminant validity (54), the internal consistency in our sample was 0.89. The Body Dissatisfaction Scale is a seven-item scale that assesses discontentment with the size and shape of various body parts that are of particular concern to those with eating disorders (e.g., the stomach). The internal consistency in our sample was 0.94.

## Neuroimaging Data Acquisition

### Image Acquisition and Preprocessing

Participants were scanned on a 3-T GE MR 750 system. T1-weighted structural scans were acquired using a 3D FSPGR BRAVO pulse sequence (repetition time, 7.58 milliseconds; echo time, 2.936 milliseconds; flip angle, 12 degrees; image matrix, 256 × 256; voxel size, 1 × 1 × 1 mm; 206 contiguous axial slices). T1-weighted images were preprocessed using the VBM8 toolbox (<http://dbm.neuro.uni-jena.de/vbm>) as implemented in SPM8 (Wellcome Department of Cognitive Neurology) to estimate the volume of gray matter, white matter, and cerebral spinal fluid for computation of total intracranial volume for the purpose of correcting insular volumes.

### Tracing

Insula volumes were traced from raw T1 images. The total volume of each insula subregion was calculated for each participant by summing across all voxels (each had a volume of .001 ml). To account for changes in overall brain volume, insula volumes were scaled relative to the total intracranial volume as computed using the VBM8 software. Inferential analyses were conducted on proportionally scaled data.

Insular volumes were manually traced using ITK-SNAP software (<http://www.itksnap.org>; (57)), dividing the insula into anterior and posterior regions as in Ref. (58). Tracing started by identifying the central insular sulcus on two parasagittal slices as a point of reference, and tracing posterior and anterior aspects of the insula separately on successive caudal-to-rostral coronal slices. In the sagittal plane, the superior circular sulcus of the insula was used to

demarcate the superior boundary of the insula and the inferior circular sulcus of the insula determined the inferior boundary. The anterior boundary of the insula was defined using the orbitoinsular sulcus, and the fusion of the superior and inferior circular sulci was used to identify the posterior boundary. Tracing proceeded rostrally from the deepest caudal point of the superior circular sulcus of the insula to the inferior circular sulcus of the insula, and orbitoinsular sulcus disappeared into the most ventral lateral point at the anterior aspect of the insula. Throughout the rostrocaudal extent of the insula, the central insular sulcus was used to segment the short anterior and long posterior insular gyri. A random subset of scans from 16 participants was scored by an additional experimenter. Consistency between volume estimates was assessed and fell into the “excellent” range (59). (See supplemental materials, Supplemental Digital Content 1, <http://links.lww.com/PSYMED/A410>, for additional validation information.)

### Statistical Methods

Given the exploratory nature of this study, we present plots of raw values and emphasize effect size estimates over statistical significance, although we do present both. The aim of the study was to (a) examine the association between insular volume and AN and (b) explore differences in that association at different points over the course of the disorder. As brain structure and size change rapidly during adolescence, age and BMI (normalized:  $z$  score) were included as covariates. Thus, study focus was on the association between MRI-derived measures of insular volume and illness parameters (duration of illness) including subjective measures of body dissatisfaction, a central psychological indicator of anorexia. Where appropriate, bivariate associations between groups were estimated and tested using both Spearman correlations (adjusted Fisher  $z$  transformation) and nonparametric rank Mann-Whitney-Wilcoxon procedures; correlational analyses designed to provide concordant tests of multivariable findings (see hereinafter), were partialled for age and BMI. Although not all tested instances failed to satisfy distributional assumptions for parametric testing, numerous instances did, leading us to apply nonparametric tests in all instances to facilitate comparisons. Bivariate investigations were expanded using multivariable regression modeling to accommodate putative covariates and modifiers as well as diagnostic status (group). The latter models included both linear and nonlinear (i.e., quadratic, piecemeal) regressions. In some instances, models were extended to examine putative differential effects for age by group using interaction terms; BMI was included as a covariate in all instances. Model fit was examined using standard diagnostics including examination of residuals, leverage, and outliers. Analyses were supplemented using graphical procedures including box-and-whisker plots, scatter plots, and loess regressions. Specifically, analyses were guided by the following hypotheses: hypothesis 1, for our investigation of age-related differences in volume between individuals with AN and HC, we hypothesized that anatomical differences will be right sided and will occur in both the anterior and posterior regions relative to healthy controls. Tests of hypothesis 1 were based on an analysis of covariance approach using a dichotomous factor denoting diagnostic status (AN = 1; HC = 0) to test for volumetric differences between groups over time. Thus, measures of insular volume were regressed on a model including age (standardized), the proxy variable denoting diagnosis, an interaction term crossing the latter two factors, and BMI (standardized); in some instances, models were reestimated after the inclusion of a quadratic term for time. Support for the hypothesis was based on the significance of the interaction term. Hypothesis 2 examined the effects of insular volume on illness duration. We hypothesized that volume size in both the right anterior and right posterior insula would correlate with illness parameters independent of BMI. Hypothesis 3 examined the association of insular volume with subjective measures of body image evaluation. We hypothesized that the volume of the right posterior insular cortex would be associated with body dissatisfaction. Tests of hypotheses 2 and 3 were restricted to the diagnostic cohort. In each instance, measures of insular volume were regressed on scale factors denoting, respectively, illness severity

and body dissatisfaction. Support for both hypotheses was based on significance of the latter terms. Models included covariates for age and BMI. For all models, we examined measures of fit (studentized residuals), influence (Cook's  $D$  and  $DF$  fits), and leverage ( $DFbetas$ )—all of which were found to be within acceptable limits. Models were estimated using SAS 9.3 statistical software.

## RESULTS

### Demographic Features

Table 1 presents demographic characteristics of the entire sample and features of the illness course and severity of the clinical sample. Average age at the time of interview was 17 years, with a range between 11 and 20 years. Groups did not differ on age or BMI. Most of the sample was white (83%). The average age of onset of AN was 13.78 years, with an average time to treatment of 11.35 months and an average length of illness of 34.64 months. Based on Mann-Whitney-Wilcoxon tests, measures of eating disorder symptoms differed significantly between groups on the Drive for Thinness and Body Dissatisfaction subscales ( $z = -3.5$  [ $p < .001$ ] and  $z = -3.6$  [ $p < .001$ ], respectively), but not the Bulimia subscale ( $z = -0.81$  [ $p > .40$ ]).

### Age

Table S2, Supplemental Digital Content 1 (<http://links.lww.com/PSYMED/A410>), presents global comparisons of white matter, gray matter, cerebrospinal fluid, and total intracranial volume (the sum of gray matter, white matter, and cerebrospinal fluid). There were no differences in these global volumes across groups. All subsequent examinations of insular volumes are the percentage of insula volumes relative to these volumes.

Results for hypothesis 1, which examined between-group differences in insular volume by region and age (normalized) and controlled for  $zBMI$ , are presented in Figure 1 and Table S3, Supplemental Digital Content 1 (<http://links.lww.com/PSYMED/A410>). Anterior regions in both hemispheres exhibited a gradual U-shaped curvilinear pattern over time, with model fit significantly improved in both instances after the addition of a quadratic (squared) term for age. Tests of main effects for group (see Table S3, Supplemental Digital Content 1, <http://links.lww.com/PSYMED/A410>, for all insular region model coefficients) and of interaction terms crossing the age components with group status were nonsignificant in the anterior region. Correlations of volume measurements between left and right anterior regions for both control and case subjects were significant (control:  $r > 0.8073$ ,  $p < .0002$ ; case:  $r > 0.80$ ,  $p < .0001$ ). In the left posterior insular region, the association between volume and age was essentially constant over time for both control and case subjects; model fit was not significantly improved by the addition of either quadratic and/or interaction terms. In contrast, patterns of associations in the right posterior insular region varied differentially between diagnostic groups. The association between volume and age in the right posterior region among control subjects was essentially flat, whereas the association in the right posterior region among examined participants increased into mid-adolescence before declining (Fig. 1). The differential association between age and volume by diagnostic group in the right posterior insula region was statistically significant ( $\beta = -0.018$  [0.008];  $t = 2.32$ ,  $p < .02$ ). The robustness of this interaction was further confirmed by examining hemispheric associations: the correlation of

**TABLE 1.** Demographic and Illness Characteristics of the Sample

	Participants With a History of AN		Typically Developing Controls		Pr >  z
	n	Mean or %	n	Mean or %	
<b>Demographic Variables</b>					
Age z score	21	0.23	20	0.16	0.86
Age, y	21	17.48	20	17.40	1.85
Asian	2	8.70	1	4.55	0.26
Black	1	4.35	4	18.18	0.50
Other	1	4.35	2	9.09	
White	19	82.61	15	68.18	
<b>Stage/Severity of Illness</b>					
BMI z score	21	-0.35	20	0.02	0.84
BMI, kg/m <sup>2</sup>	21	20.01	20	21.34	3.00
Body dissatisfaction <sup>a</sup>	21	15.15	17	5.47	5.39
Drive for thinness <sup>a</sup>	21	15.40	17	4.82	6.07
Bulimia	20	3.70	17	1.59	1.70
Age of onset, y	21	13.78	n/a	n/a	n/a
Duration of illness <sup>b</sup>	20	3.14	n/a	n/a	n/a
Length of weight restoration <sup>b</sup>	21	3.33	n/a	n/a	n/a
<b>Length of Weight Restoration by Category<sup>c</sup> (n = 21)</b>					
Unhealthy weight	5	24			
<3 mo	1	4.8			
3–6 mo	6	28.6			
7–9 mo	1	4.8			
10–12 mo	0	0			
Over a year	2	9.5			
1–2 y	1	4.8			
>2 y	5	23.8			
<b>Drive for Thinness Classification (Weight-Restored Group, n = 16)</b>					
Clinical range	10	63			
Normative range	5	31			
Missing	1	6			
Lowest BMI while ill	21	15.19			2.53

Group Membership	n	% of Group	Control (n)	% of Group	AN (n)	% of Group
Current AN	5	23.81	1	5	4	24
Weight-restored AN	16	76.12	0	0	14	67
History of purging	3	21.74				
<b>Model Covariates</b>						
Left-handed						
Psychotropic medication						

AN = anorexia nervosa; SD = standard deviation; BMI = body mass index.

<sup>a</sup>Subscale scores from the Eating Disorder Inventory (Third Edition).

<sup>b</sup>Number and proportion of years.

<sup>c</sup>At the time of intake interview.

volume measurements between left and right posterior insular regions for control subjects was statistically significant ( $r = 0.75, p < .001$ ). In contrast, the left/right correlation of volumes in right posterior regions for clinical participants did not differ significantly from zero ( $r = 0.04, p > .80$ ); that is, volumes from the left and right posterior cortex were unrelated in clinical participants. Associations between insular volumes and age of onset, handedness, and medication status were tested both individually and concurrently; in all instances, associations were nonsignificant (data not shown). Thus, in support of our initial hypothesis, volume differences were noted in the right posterior insular cortex in AN participants relative to the control group. In contrast to initial hypotheses, differences between volumes in anterior regions did not differ by group.

### Illness Severity

The tests of our second hypothesis positing associations between volumetric abnormalities in insular regions and indices of illness severity are presented in Table 2 and Figure 2 (model coefficients are in Table S4, Supplemental Digital Content 1, <http://links.lww.com/PSYMED/A410>). In anterior insular regions, residual volumes (adjusted for normalized measures of age and BMI) decreased with length of illness to a midpoint of approximately 40 months where, after the association, became constant. The magnitude of the decrease was of large effect and statistically significant in the right anterior region ( $r = -0.5, p < .03$ ). Although the loess fit (Fig. 2) suggested a nonlinear association between length of illness and volume, adding a quadratic component to the models did not significantly improve fit (data not shown). In contrast, the association in the right posterior region exhibited a pronounced positive association—marked by an almost constant increase with increasing length of illness ( $r = 0.5, p < .04$ ). The correlation in the left posterior region was essentially zero.


### Body Dissatisfaction

The test of our third hypothesis examined the associations between insular volumes and a subjective measure of body dissatisfaction (Eating Disorder Inventory Body Dissatisfaction Subscale; Table 2). Findings replicated the earlier pattern of results: a negative association between volume measurements in the right anterior insula and the index of body dissatisfaction ( $r = -0.49, p < .04$ ) and a positive association between volume measurements in the right posterior insula and the index of dissatisfaction ( $r = 0.40, p < .10$ ; Table 2).

### DISCUSSION

The most robust and consistent finding across all research questions was related to the volume of the right posterior insula cortex. Our results show (a) age-related differences relative to healthy controls, (b) a positive correlation with duration of illness, and (c) a positive association with a subjective measure of body dissatisfaction, although this moderately sized effect only approached statistical significance. This distinct pattern contrasted with those observed in either bilateral anterior regions of the insula or the left posterior region (Figs. 1, 2).

Our data suggest that increased right posterior insular volume may be worthy of further exploration as a putative marker of illness progression or as a vulnerability factor for severity of illness in AN. The region receives and encodes visceral interoceptive input relayed



**FIGURE 1.** Age by posterior and anterior insular volume in clinical participants relative to controls. Volume size as a function of age and group membership, controlling for body mass index, is depicted. Reported volumes are the proportion of two volumes ml (insula)/ml (ICV). Age was standardized and thus the  $x$ -axis reflects  $z$  scores that reflect relative age within the sample. The dashed line depicts adolescents with a current or past diagnosis of AN ( $n = 21$ ); the solid line depicts typically developing adolescent control participants ( $n = 20$ ). Circles represent healthy controls; triangles, current diagnosis of AN; square, history of AN, but weight restored. Tests of main and interaction effects in bilateral anterior regions and the left posterior were nonsignificant. The differential association between age and volume by diagnostic group in the right posterior insula region was statistically significant ( $\beta = -0.018$  [0.008];  $t = 2.32$ ,  $p = .02$ ). ICV = intracranial volume; AN = anorexia nervosa. Color image is available only in online version ([www.psychosomaticmedicine.org](http://www.psychosomaticmedicine.org)).

via the nuclei in the solitary tract and thalamus (60). How structural irregularities in a neural region that relays somatosensory and pain information are putatively associated with illness pathophysiology in AN is unclear: our cross-sectional data cannot adjudicate how the structure of the posterior insular cortex is associated with illness pathophysiology. Notwithstanding, consideration of various hypothetical scenarios can inform the design of future studies that can address if and how afferent visceral input relates to the core pathology of AN.

For instance, one hypothesis is that the pathophysiology of AN is partially caused by or associated with developmental alterations in afferent signaling from the viscera. Given the early maturation of this region (e.g., see Mayer (23) for review), one could postulate that early life events that sensitize pathways of the gut-brain axis may influence the development of this structure. In support of this, there is evidence of an excess of early gastrointestinal events, eating difficulties, and somatic symptoms before illness onset in AN (19–22). Notably, although some gastrointestinal symptoms such as abdominal pain persist beyond weight restoration, many gastrointestinal symptoms also arise as a function of eating

disorder symptoms and complicate treatment (61–63). Such signaling is mediated by the vagus nerve, and atypical vagal tone has also been reported in AN (although the trajectory of development in relation to illness course is unclear) (64). Thus, one possibility is that early experiences that sensitize pain pathways may influence disorder emergence via alterations in insula morphology that may be further altered by AN.

A related hypothesis is that the context of starvation itself influences the structure of this region. Notably, AN has been consistently associated with global and focal reductions of gray matter volume, with some evidence documenting a restoration and normalization of cortical mass with weight restoration (42–44,65). Findings of *increased* volume are sparse but noteworthy (41,42). Frank et al. (41) reported, as we did, increased volume in the right insular cortex in adolescents with AN. Using voxel-based morphometry, peak differences were localized to a region more anterior than those in the current study, just anterior to the central sulcus in the mid-insula. Amianto et al. (42) reported increased volume of the somatosensory cortex in adults with AN relative

**TABLE 2.** Association of Insular Volume With Illness Parameters

Insular Region	Pearson <i>r</i>	Fisher <i>z</i>	-95 CI	+95 CI	<i>p</i> H 0: $\rho = 0$
Length of illness <sup>a</sup>					
Left anterior	-0.38	-0.40	-0.72	0.11	0.13
Right anterior	-0.50	-0.55	-0.78	-0.05	0.03
Left posterior	-0.05	-0.05	-0.51	0.42	0.83
Right posterior	0.48	0.52	0.02	0.77	0.04
Body dissatisfaction <sup>b</sup>					
Left anterior	-0.38	-0.40	-0.72	0.11	0.12
Right anterior	-0.49	-0.53	-0.78	-0.02	0.04
Left posterior	-0.38	-0.40	-0.72	0.11	0.12
Right posterior	0.40	0.43	-0.08	0.73	0.10

CI = confidence interval.


<sup>a</sup>Length measures in years and proportion of years.

<sup>b</sup>Body dissatisfaction subscale of the Eating Disorder Order Inventory.

to controls. Although these structural abnormalities cannot point to specific pathways of disorder pathophysiology, these findings do highlight the importance of further study of regions such as the somatosensory cortex and associated somatomotor regions to inform the nature of experience in AN.

Related to this hypothesis is that starvation and associated disruptions in hormone signaling may delay typical neurodevelopmental changes. Thus, current findings may be reflections of this delay. Dennis et al. (39) investigated the functional connectivity of insular networks across puberty and adulthood. Results revealed that although connections largely decreased in anterior insular regions, these connections increased in posterior insular-temporal regions (39). Furthermore, Takahashi et al. (66) reported that bilateral insular gray matter volume is negatively correlated with age in a sample of adults (ages,  $\geq 20$  years). Better understanding of the effects of starvation at specific developmental stages may help to elucidate neurodevelopmental factors that maintain the disorder.

However, to understand the role of atypical structure in body image disturbance in AN, novel designs are needed. Our data demonstrated a moderate (statistically insignificant) association between



**FIGURE 2.** The relationship between length of illness and insular volume in adolescents with AN. The relationship between length of illness in months and volume in anterior and posterior insular components in 21 adolescents with a current or previous diagnosis of AN is depicted. Reported volumes are the proportion of two volumes ml (insula)/ml (ICV). The magnitude of association was of large effect but in opposite direction: being negatively associated with right anterior insular volume ( $r = -0.5, p = .03$ ) and positively associated with right posterior volume ( $r = 0.5, p = .04$ ). Triangle represents current diagnosis of AN; square, history of AN, but weight restored. AN = anorexia nervosa; ICV = intracranial volume. Color image is available only in online version ([www.psychosomaticmedicine.org](http://www.psychosomaticmedicine.org)).

size of the right posterior insular cortex and body dissatisfaction and the inverse association with the right anterior insula and body dissatisfaction. However, to elucidate the significance of brain structure, brain function, and the mediators of the structure-function relationship, there are several links in the mechanistic chain missing from this investigation. Studies that characterize structure, the efficiency of functional circuits mediating afferent signaling and those circuits that mediate awareness and interpretation of that signaling, and manipulations of state conditions that can inform how features of AN may alter interoceptive experience are needed. As described in the work by Herbert et al. (67), interoceptive signals are sensed, interpreted, and then evaluated. Each of these processes can independently contribute to eating behavior and body experience.

Previous work in typically developing controls indicates that behaviors associated with AN impact interoceptive sensitivity, and that sensitivity to interoceptive signals and the interpretation of those signals are independent processes that differentially impact eating behavior. For example, Herbert et al. (67) investigated the effects of an acute fast on interoceptive sensitivity, finding that sensitivity to heartbeat and experienced hunger increases after an acute fast, partially due to changes in cardiodynamic activity. Furthermore, Herbert et al. (26) reported that interoceptive sensitivity and evaluations of the aversiveness of interoceptive experience independently contributed to eating behavior. Exercise is an additional example of a behavior associated with AN that may alter the strength of interoceptive signaling and the perception of those signals. Exercise has also been shown to increase interoceptive awareness and accuracy, highlighting a functional path that may correspond with possible changes in structure. As noted, exercise is associated with alternations in brain-derived neurotrophic factor function, a mechanism whereby the symptoms of AN may alter brain structure. Combined, this pattern of findings points to how aberrations in lower level encoding, when combined with behaviors and attitudes that may influence interoceptive experience, may theoretically contribute to body image disturbance in AN. Thus, one hypothetical model of the pathophysiology of anorexia is that individuals who are premorbidly viscerally hypersensitive (as due to experiences that sensitize gut brain pathways) may be differentially or more strongly reinforced by events/behaviors that alter interoceptive experience, and this may influence the interpretation and meaning of these experiences.

Furthermore, the behaviors that constitute the illness of AN may differentially influence these processes over time, and thus, the model of interoceptive sensitivity and accuracy may alter throughout the illness of AN. Thus, future studies need to incorporate behavioral tasks that assess interoceptive sensitivity, the interpretation and evaluation of interoceptive signals, and the functional circuitry in structural architecture that support these processes. Finally, given our findings regarding a region implicated in somatomotor processing, future work that particularly emphasizes the somatosensory cortex would be of special interest.

This study had several limitations. The clinical significance of findings from our cross-sectional design requires verification with a longitudinal design. Although our use of population screening methods with limited exclusion criteria strengthens generalizability, it weakens internal validity because many participants were on psychotropic medications. Inclusion of a medication proxy did not alter

study findings, and given the limited empirical evidence regarding the effectiveness of a psychotropic agent on AN symptoms, the concern is lessened. Finally, our sample size, although typical of published neuroimaging studies, was small and so precluded the detection of smaller but still clinically significant effects.

In summary, this evidence highlights the need for future work to explore a new model of body image disturbance that further examines the role of sensory experience. This model may lead to novel conceptualizations of AN and avenues for intervention such as those that help those with AN to experience and contextualize volatile somatic signals to better guide adaptive behavior. Such an approach might offer an alternative or supplement to cognitively focused interventions and enhance treatment effectiveness (68).

*Source of Funding and Conflicts of Interest:* This work was supported by the National Institute of Mental Health (Grant R01-MH-088678). The authors have no conflict of interest in regard to this article.

## REFERENCES

- Halmi KA. Perplexities of treatment resistance in eating disorders. *BMC Psychiatry* 2013;13:292.
- Grunwald M, Ettrich C, Krause W, Assmann B, Dahne A, Weiss T, Gertz HJ. Haptic perception in anorexia nervosa before and after weight gain. *J Clin Exp Neuropsychol* 2001;23:520–9.
- Arcelus J, Mitchell AJ, Wales J, Nielsen S. Mortality rates in patients with anorexia nervosa and other eating disorders: a meta-analysis of 36 studies. *Arch Gen Psychiatry* 2011;68:724–31.
- Eisler I, Simic M, Hodsoll J, Asen E, Berelowitz M, Connan F, Ellis G, Hugo P, Schmidt U, Treasure J, Yi I, Landau S. A pragmatic randomised multi-centre trial of multifamily and single family therapy for adolescent anorexia nervosa. *BMC Psychiatry* 2016;16:422.
- Le Grange D, Hughes EK, Court A, Yeo M, Crosby RD, Sawyer SM. Randomized clinical trial of parent-focused treatment and family-based treatment for adolescent anorexia nervosa. *J Am Acad Child Adolesc Psychiatry* 2016;55:683–92.
- Agras WS, Lock J, Brandt H, Bryson SW, Dodge E, Halmi KA, Jo B, Johnson C, Kaye W, Wilfley D, Woodside B. Comparison of 2 family therapies for adolescent anorexia nervosa: a randomized parallel trial. *JAMA Psychiatry* 2014;71:1279–86.
- Grunwald M, Ettrich C, Assmann B, Dähne A, Krause W, Busse F, Gertz HJ. Deficits in haptic perception and right parietal theta power changes in patients with anorexia nervosa before and after weight gain. *Int J Eat Disord* 2001;29:417–28.
- Halmi KA, Agras WS, Crow S, Mitchell J, Wilson GT, Bryson SW, Kraemer HC. Predictors of treatment acceptance and completion in anorexia nervosa: implications for future study designs. *Arch Gen Psychiatry* 2005;62:776–81.
- Towell DB, Woodford S, Reid S, Rooney B, Towell A. Compliance and outcome in treatment-resistant anorexia and bulimia: a retrospective study. *Br J Clin Psychol* 2001;40:189–95.
- Cash TF, Deagle EA 3rd. The nature and extent of body-image disturbances in anorexia nervosa and bulimia nervosa: a meta-analysis. *Int J Eat Disord* 1997; 22:107–25.
- Gardner RM, Brown DL. Body size estimation in anorexia nervosa: a brief review of findings from 2003 through 2013. *Psychiatry Res* 2014;219:407–10.
- Gaudio S, Brooks SJ, Riva G. Nonvisual multisensory impairment of body perception in anorexia nervosa: a systematic review of neuropsychological studies. *PLoS One* 2014;9:e110087.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed. Washington, DC: American Psychiatric Association; 2013.
- Pollatos O, Kurz AL, Albrecht J, Schredler T, Kleemann AM, Schöpf V, Kopietz R, Wiesmann M, Schandry R. Reduced perception of bodily signals in anorexia nervosa. *Eat Behav* 2008;9:381–8.
- Gupta MA, Gupta AK, Schork NJ, Wateel GN. Perceived touch deprivation and body image—some observations among eating-disordered and nonclinical subjects. *J Psychosom Res* 1995;39:459–64.
- Keizer A, Smeets MA, Dijkerman HC, Uzunbajakau SA, van Elburg A, Postma A. Too fat to fit through the door: first evidence for disturbed body-scaled action in anorexia nervosa during locomotion. *PLoS One* 2013;8:e64602.
- Zucker NL, Merwin RM, Bulik CM, Moskovich A, Wildes JE, Groh J. Subjective experience of sensation in anorexia nervosa. *Behav Res Ther* 2013;51: 256–65.

18. Merwin RM, Moskovich AA, Wagner HR, Ritschel LA, Craighead LW, Zucker NL. Emotion regulation difficulties in anorexia nervosa: relationship to self-perceived sensory sensitivity. *Cogn Emot* 2013;27:441–52.
19. Jacobi C, Hayward C, de Zwaan M, Kraemer HC, Agras WS. Coming to terms with risk factors for eating disorders: application of risk terminology and suggestions for a general taxonomy. *Psychol Bull* 2004;130:19–65.
20. Marchi M, Cohen P. Early childhood eating behaviors and adolescent eating disorders. *J Am Acad Child Adolesc Psychiatry* 1990;29:112–7.
21. Kotler LA, Cohen P, Davies M, Pine DS, Walsh BT. Longitudinal relationships between childhood, adolescent, and adult eating disorders. *J Am Acad Child Adolesc Psychiatry* 2001;40:1434–40.
22. Silberg JL, Bulik CM. The developmental association between eating disorders symptoms and symptoms of depression and anxiety in juvenile twin girls. *J Child Psychol Psychiatry* 2005;46:1317–26.
23. Mayer EA. Gut feelings: the emerging biology of gut-brain communication. *Nat Rev Neurosci* 2011;12:453–66.
24. Zucker NL, Harshaw C. Emotion, attention, and relationships: a developmental model of self-regulation in anorexia nervosa and related disordered eating behaviors. In: Lock J, editor. *The Oxford Handbook of Developmental Perspectives on Child and Adolescent Eating Disorders*. London: Oxford Press; 2012:67–87.
25. Bechara A, Damasio AR. The somatic marker hypothesis: a neural theory of economic decision. *Games Econ Behav* 2005;52:336–72.
26. Herbert BM, Blechert J, Hautzinger M, Matthias E, Herbert C. Intuitive eating is associated with interoceptive sensitivity. Effects on body mass index. *Appetite* 2013;70:22–30.
27. Kaye WH, Fudge JL, Paulus M. New insights into symptoms and neurocircuit function of anorexia nervosa. *Nat Rev Neurosci* 2009;10:573–84.
28. Craig AD. How do you feel—now? The anterior insula and human awareness. *Nat Rev Neurosci* 2009;10:59–70.
29. Kurth F, Zilles K, Fox PT, Laird AR, Eickhoff SB. A link between the systems: functional differentiation and integration within the human insula revealed by meta-analysis. *Brain Struct Funct* 2010;214:519–34.
30. Chang LJ, Yarkoni T, Khaw MW, Sanfey AG. Decoding the role of the insula in human cognition: functional parcellation and large-scale reverse inference. *Cereb Cortex* 2013;23:739–49.
31. Pollatos O, Kirsch W, Schandry R. Brain structures involved in interoceptive awareness and cardioafferent signal processing: a dipole source localization study. *Hum Brain Mapp* 2005;26:54–64.
32. Droutman V, Bechara A, Read SJ. Roles of the different sub-regions of the insular cortex in various phases of the decision-making process. *Front Behav Neurosci* 2015;9:309.
33. Yu YB, Zuo XL, Zhao QJ, Chen FX, Yang J, Dong YY, Wang P, Li YQ. Brain-derived neurotrophic factor contributes to abdominal pain in irritable bowel syndrome. *Gut* 2012;61:685–94.
34. Church DD, Hoffman JR, Mangine GT, Jajtner AR, Townsend JR, Beyer KS, Wang R, La Monica MB, Fukuda DH, Stout JR. Comparison of high-intensity vs. high-volume resistance training on the BDNF response to exercise. *J Appl Physiol* (1985) 2016;121:123–8.
35. Gogtay N, Giedd JN, Lusk L, Hayashi KM, Greenstein D, Vaituzis AC, Nugent TF 3rd, Herman DH, Clasen LS, Toga AW, Rapoport JL, Thompson PM. Dynamic mapping of human cortical development during childhood through early adulthood. *Proc Natl Acad Sci U S A* 2004;101:8174–9.
36. Prado EL, Dewey KG. Nutrition and brain development in early life. *Nutr Rev* 2014;72:267–84.
37. Raznahan A, Shaw PW, Lerch JP, Clasen LS, Greenstein D, Berman R, Pipitone J, Chakravarty MM, Giedd JN. Longitudinal four-dimensional mapping of subcortical anatomy in human development. *Proc Natl Acad Sci U S A* 2014;111:1592–7.
38. Kurth F, Eickhoff SB, Schleicher A, Hoemke L, Zilles K, Amunts K. Cytoarchitecture and probabilistic maps of the human posterior insular cortex. *Cereb Cortex* 2010;20:1448–61.
39. Dennis EL, Jahanshad N, McMahon KL, de Zubicaray GI, Martin NG, Hickie IB, Toga AW, Wright MJ, Thompson PM. Development of insula connectivity between ages 12 and 30 revealed by high angular resolution diffusion imaging. *Hum Brain Mapp* 2014;35:1790–800.
40. Merwin RM, Zucker NL, Lacy JL, Elliott CA. Interoceptive awareness in eating disorders: distinguishing lack of clarity from non-acceptance of internal experience. *Cognit Emot* 2010;24:892–902.
41. Frank GK, Shott ME, Hagman JO, Yang TT. Localized brain volume and white matter integrity alterations in adolescent anorexia nervosa. *J Am Acad Child Adolesc Psychiatry* 2013;52:1066–75.
42. Amianto F, Caroppo P, D'Agata F, Spalatro A, Lavagnino L, Caglio M, Righi D, Bergui M, Abbate-Daga G, Rigardetto R, Mortara P, Fassino S. Brain volumetric abnormalities in patients with anorexia and bulimia nervosa: a voxel-based morphometry study. *Psychiatry Res* 2013;213:210–6.
43. Friederich HC, Walther S, Bendszus M, Biller A, Thomann P, Zeigermann S, Katus T, Brunner R, Zastrow A, Herzog W. Grey matter abnormalities within cortico-limbic-striatal circuits in acute and weight-restored anorexia nervosa patients. *Neuroimage* 2012;59:1106–13.
44. Gaudio S, Nocchi F, Franchin T, Genovese E, Cannata V, Longo D, Fariello G. Gray matter decrease distribution in the early stages of anorexia nervosa restrictive type in adolescents. *Psychiatry Res* 2011;191:24–30.
45. Castro-Fornieles J, Bargallo N, Lazaro L, Andres S, Falcon C, Plana MT, Junque C. A cross-sectional and follow-up voxel-based morphometric MRI study in adolescent anorexia nervosa. *J Psychiatr Res* 2008;43:331–40.
46. Joos A, Klöppel S, Hartmann A, Glauche V, Tüscher O, Perlov E, Saum B, Freyer T, Zeeck A, Tebartz van Elst L. Voxel-based morphometry in eating disorders: correlation of psychopathology with grey matter volume. *Psychiatry Res* 2010;182:146–51.
47. King JA, Geisler D, Ritschel F, Boehm I, Seidel M, Roschinski B, Soltwedel L, Zwiip J, Pfuhl G, Marxen M, Roessner V, Ehrlich S. Global cortical thinning in acute anorexia nervosa normalizes following long-term weight restoration. *Biol Psychiatry* 2015;77:624–32.
48. Suchan B, Busch M, Schulte D, Gronermeier D, Herpertz S, Vocks S. Reduction of gray matter density in the extrastriate body area in women with anorexia nervosa. *Behav Brain Res* 2010;206:63–7.
49. Giedd JN. Structural magnetic resonance imaging of the adolescent brain. *Ann N Y Acad Sci* 2004;1021:77–85.
50. Mills KL, Goddings AL, Clasen LS, Giedd JN, Blakemore SJ. The developmental mismatch in structural brain maturation during adolescence. *Dev Neurosci* 2014;36:147–60.
51. Kragel PA, Zucker NL, Covington VE, LaBar KS. Developmental trajectories of cortical-subcortical interactions underlying the evaluation of trust in adolescence. *Soc Cogn Affect Neurosci* 2015;10:240–7.
52. Li D, Zucker NL, Kragel PA, Covington VE, LaBar KS. Adolescent development of insula-dependent interoceptive regulation. *Dev Sci* 2016.
53. Franz L, Angold A, Copeland W, Costello EJ, Towse-Goodman N, Egger H. Pre-school anxiety disorders in pediatric primary care: prevalence and comorbidity. *J Am Acad Child Adolesc Psychiatry* 2013;52:1294–303.
54. Garner DM. *Eating Disorder Inventory-3: Professional Manual*, Lutz, FL: Psychological Assessment Resources, Inc.; 2004.
55. Hagman J, Gardner RM, Brown DL, Gralla J, Fier JM, Frank GK. Body size overestimation and its association with body mass index, body dissatisfaction, and drive for thinness in anorexia nervosa. *Eat Weight Disord* 2015;20:449–55.
56. Peñas-Lledó E, Bulik CM, Lichtenstein P, Larsson H, Baker JH. Risk for self-reported anorexia or bulimia nervosa based on drive for thinness and negative affect clusters/dimensions during adolescence: a three-year prospective study of the TChAD cohort. *Int J Eat Disord* 2015;48:692–9.
57. Yushkevich PA, Piven J, Hazlett HC, Smith RG, Ho S, Gee JC, Gerig G. User-guided 3D active contour segmentation of anatomical structures: significantly improved efficiency and reliability. *Neuroimage* 2006;31:1116–28.
58. Takahashi T, Suzuki M, Zhou SY, Hagino H, Tanino R, Kawasaki Y, Nohara S, Yamashita I, Seto H, Kurachi M. Volumetric MRI study of the short and long insular cortices in schizophrenia spectrum disorders. *Psychiatry Res* 2005;138:209–20.
59. Cicchetti DV. Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychol Assess* 1994;6:284.
60. Wittmann M, Simmons AN, Aron JL, Paulus MP. Accumulation of neural activity in the posterior insula encodes the passage of time. *Neuropsychologia* 2010;48:3110–20.
61. Wang XJ, Luscombe GM, Boyd C, Kellow J, Abraham S. Functional gastrointestinal disorders in eating disorder patients: altered distribution and predictors using ROME III compared to ROME II criteria. *World J Gastroenterol* 2014;20:16293–9.
62. Perkins SJ, Keville S, Schmidt U, Chalder T. Eating disorders and irritable bowel syndrome: is there a link? *J Psychosom Res* 2006;59:57–64.
63. Salvioli B, Pellicciari A, Iero L, Di Pietro E, Moscano F, Gualandi S, Stanghellini V, De Giorgio R, Ruggeri E, Franzoni E. Audit of digestive complaints and psychopathological traits in patients with eating disorders: a prospective study. *Dig Liver Dis* 2013;45:639–44.
64. Kollai M, Bonyhay I, Jokkel G, Szonyi L. Cardiac vagal hyperactivity in adolescent anorexia nervosa. *Eur Heart J* 1994;15:1113–8.
65. Boghi A, Sterpone S, Sales S, D'Agata F, Bradac GB, Zullo G, Munno D. In vivo evidence of global and focal brain alterations in anorexia nervosa. *Psychiatry Res* 2011;192:154–9.
66. Takahashi R, Ishii K, Kakigi T, Yokoyama K. Gender and age differences in normal adult human brain: voxel-based morphometric study. *Hum Brain Mapp* 2011;32:1050–8.
67. Herbert BM, Herbert C, Pollatos O, Weimer K, Enck P, Sauer H, Zipfel S. Effects of short-term food deprivation on interoceptive awareness, feelings and autonomic cardiac activity. *Biol Psychol* 2012;89:71–9.
68. Bulik CM, Berkman ND, Brownley KA, Sedway JA, Lohr KN. Anorexia nervosa treatment: a systematic review of randomized controlled trials. *Int J Eat Disord* 2007;40:310–20.